

References for

Central Alpha Agonists, Gabapentinoids, and Sodium Channel Antagonists

Nadia Awad

Central Alpha Agonists

Giovannitti JA, Thoms SM, Crawford JJ. Alpha-2 adrenergic receptor agonists: a review of current clinical applications. *Anesth Prog.* 2015;62(1):31-9.

Habib AS, Gan TJ. Role of analgesic adjuncts in postoperative pain management. *Anesthesiol Clin North America.* 2005;23(1):85-107.

Zhu QM, Lesnick JD, Jasper JR, et al. Cardiovascular effects of rilmenidine, moxonidine and clonidine in conscious wild-type and D79N alpha2A-adrenoceptor transgenic mice. *Br J Pharmacol.* 1999;126(6):1522-30.

Hall DL, Rezvan E, Tatakis DN, Walters JD. Oral clonidine pretreatment prior to venous cannulation. *Anesth Prog.* 2006;53(2):34-42.

Smith H, Elliott J. Alpha(2) receptors and agonists in pain management. *Curr Opin Anaesthesiol.* 2001;14(5):513-8.

Pal SK, Cortiella J, Herndon D. Adjunctive methods of pain control in burns. *Burns.* 1997;23(5):404-12.

Virtanen R, Savola JM, Saano V, Nyman L. Characterization of the selectivity, specificity and potency of medetomidine as an alpha 2-adrenoceptor agonist. *Eur J Pharmacol.* 1988;150(1-2):9-14.

Jaakola ML, Salonen M, Lehtinen R, Scheinin H. The analgesic action of dexmedetomidine--a novel alpha 2-adrenoceptor agonist--in healthy volunteers. *Pain.* 1991;46(3):281-5.

Cao J, Shi X, Miao X, Xu J. Effects of premedication of midazolam or clonidine on perioperative anxiety and pain in children. *Biosci Trends.* 2009;3(3):115-8.

Yanagidate F, Hamaya Y, Dohi S. Clonidine premedication reduces maternal requirement for intravenous morphine after cesarean delivery without affecting newborn's outcome. *Reg Anesth Pain Med.* 2001;26(5):461-7.

Marinangeli F, Ciccozzi A, Donatelli F, et al. Clonidine for treatment of postoperative pain: a dose-finding study. *Eur J Pain*. 2002;6(1):35-42.

Bergendahl HT, Lönnqvist PA, Eksborg S, et al. Clonidine vs. midazolam as premedication in children undergoing adeno-tonsillectomy: a prospective, randomized, controlled clinical trial. *Acta Anaesthesiol Scand*. 2004;48(10):1292-300.

Blaudszun G, Lysakowski C, Elia N, Tramèr MR. Effect of perioperative systemic $\alpha 2$ agonists on postoperative morphine consumption and pain intensity: systematic review and meta-analysis of randomized controlled trials. *Anesthesiology*. 2012;116(6):1312-22.

Sung CS, Lin SH, Chan KH, Chang WK, Chow LH, Lee TY. Effect of oral clonidine premedication on perioperative hemodynamic response and postoperative analgesic requirement for patients undergoing laparoscopic cholecystectomy. *Acta Anaesthesiol Sin*. 2000;38(1):23-9.

Lyons B, Casey W, Doherty P, Mchugh M, Moore KP. Pain relief with low-dose intravenous clonidine in a child with severe burns. *Intensive Care Med*. 1996;22(3):249-51.

Giovannoni MP, Ghelardini C, Vergelli C, Dal Piaz V. Alpha2-agonists as analgesic agents. *Med Res Rev*. 2009;29(2):339-68.

Deer T, Krames ES, Hassenbusch SJ, et al. Polyanalgesic consensus conference 2007: recommendations for the management of pain by intrathecal (intraspinial) drug delivery: report of an interdisciplinary expert panel. *Neuromodulation*. 2007;10(4):300-28.

Filos KS, Goudas LC, Patroni O, Polyzou V. Hemodynamic and analgesic profile after intrathecal clonidine in humans. A dose-response study. *Anesthesiology*. 1994;81(3):591-601.

Axelsson K, Gupta A. Local anaesthetic adjuvants: neuraxial versus peripheral nerve block. *Curr Opin Anaesthesiol*. 2009;22(5):649-54.

De Kock MF, Pichon G, Scholtes JL. Intraoperative clonidine enhances postoperative morphine patient-controlled analgesia. *Can J Anaesth*. 1992;39(6):537-44.

Huang YS, Lin LC, Huh BK, et al. Epidural clonidine for postoperative pain after total knee arthroplasty: a dose-response study. *Anesth Analg*. 2007;104(5):1230-5.

Förster JG, Rosenberg PH. Small dose of clonidine mixed with low-dose ropivacaine and fentanyl for epidural analgesia after total knee arthroplasty. *Br J Anaesth*. 2004;93(5):670-7.

Eisenach JC, Dupen S, Dubois M, Miguel R, Allin D. Epidural clonidine analgesia for

intractable cancer pain. The Epidural Clonidine Study Group. *Pain*. 1995;61(3):391-9.

McCartney CJ, Duggan E, Apatu E. Should we add clonidine to local anesthetic for peripheral nerve blockade? A qualitative systematic review of the literature. *Reg Anesth Pain Med*. 2007;32(4):330-8.

Pöpping DM, Elia N, Marret E, Wenk M, Tramèr MR. Clonidine as an adjuvant to local anesthetics for peripheral nerve and plexus blocks: a meta-analysis of randomized trials. *Anesthesiology*. 2009;111(2):406-15.

Wrzosek A, Woron J, Dobrogowski J, Jakowicka-wordliczek J, Wordliczek J. Topical clonidine for neuropathic pain. *Cochrane Database Syst Rev*. 2015;8:CD010967.

Venn RM, Karol MD, Grounds RM. Pharmacokinetics of dexmedetomidine infusions for sedation of postoperative patients requiring intensive care. *Br J Anaesth*. 2002;88(5):669-75.

Wong A, Smithburger PL, Kane-gill SL. Review of adjunctive dexmedetomidine in the management of severe acute alcohol withdrawal syndrome. *Am J Drug Alcohol Abuse*. 2015 [Epub ahead of print]

Bielka K, Kuchyn I, Glumcher F. Addition of dexmedetomidine to benzodiazepines for patients with alcohol withdrawal syndrome in the intensive care unit: a randomized controlled study. *Ann Intensive Care*. 2015;5(1):33.

Vanderweide LA, Foster CJ, Maclaren R, Kiser TH, Fish DN, Mueller SW. Evaluation of Early Dexmedetomidine Addition to the Standard of Care for Severe Alcohol Withdrawal in the ICU: A Retrospective Controlled Cohort Study. *J Intensive Care Med*. 2016;31(3):198-204.

Beg M, Fisher S, Siu D, Rajan S, Troxell L, Liu VX. Treatment of Alcohol Withdrawal Syndrome with and without Dexmedetomidine. *Perm J*. 2016;20(2):49-53.

Hayes BD, Nelson LS. Letter in Response to Evaluation of Early Dexmedetomidine Addition to the Standard of Care for Severe Alcohol Withdrawal in the ICU: A Retrospective Controlled Cohort Study. *J Intensive Care Med*. 2016;31(5):353-4.

Nichols DP, Berkenbosch JW, Tobias JD. Rescue sedation with dexmedetomidine for diagnostic imaging: a preliminary report. *Paediatr Anaesth*. 2005;15(3):199-203.

Berkenbosch JW, Wankum PC, Tobias JD. Prospective evaluation of dexmedetomidine for noninvasive procedural sedation in children. *Pediatr Crit Care Med*. 2005;6(4):435-9.

Koroglu A, Demirbilek S, Teksan H, Sagir O, But AK, Ersoy MO. Sedative, haemodynamic and respiratory effects of dexmedetomidine in children undergoing magnetic resonance imaging examination: preliminary results. *Br J Anaesth*. 2005;94(6):821-4.

Mason KP, Zgleszewski SE, Prescilla R, Fontaine PJ, Zurakowski D. Hemodynamic effects of dexmedetomidine sedation for CT imaging studies. *Paediatr Anaesth*. 2008;18(5):393-402.

Jewett J, Phillips WJ. Dexmedetomidine for procedural sedation in the emergency department. *Eur J Emerg Med*. 2010;17(1):60.

Godwin SA, Burton JH, Gerardo CJ, et al. Clinical policy: procedural sedation and analgesia in the emergency department. *Ann Emerg Med*. 2014;63(2):247-58.e18.

Tobias JD. Dexmedetomidine and ketamine: an effective alternative for procedural sedation?. *Pediatr Crit Care Med*. 2012;13(4):423-7.

Goyal R, Singh S, Shukla RN, Patra AK, Bhargava DV. Ketodex, a combination of dexmedetomidine and ketamine for upper gastrointestinal endoscopy in children: a preliminary report. *J Anesth*. 2013;27(3):461-3.

Kako H, Corridore M, Kean J, Mendell JR, Flanigan KM, Tobias JD. Dexmedetomidine and ketamine sedation for muscle biopsies in patients with Duchenne muscular dystrophy. *Paediatr Anaesth*. 2014;24(8):851-6.

Goyal R, Hasnain S, Mittal S, Shreevastava S. A randomized, controlled trial to compare the efficacy and safety profile of a dexmedetomidine-ketamine combination with a propofol-fentanyl combination for ERCP. *Gastrointest Endosc*. 2016;83(5):928-33.

Hadi SM, Saleh AJ, Tang YZ, Daoud A, Mei X, Ouyang W. The effect of KETODEX on the incidence and severity of emergence agitation in children undergoing adenotonsillectomy using sevoflurane based-anesthesia. *Int J Pediatr Otorhinolaryngol*. 2015;79(5):671-6.

Ibacache ME, Muñoz HR, Fuentes R, Cortínez LI. Dexmedetomidine-ketamine combination and caudal block for superficial lower abdominal and genital surgery in children. *Paediatr Anaesth*. 2015;25(5):499-505.

Bhana N, Goa KL, McClellan KJ. Dexmedetomidine. *Drugs*. 2000;59(2):263-8.

Alhashemi JA, Kaki AM. Dexmedetomidine in combination with morphine PCA provides superior analgesia for shockwave lithotripsy. *Can J Anaesth*. 2004;51(4):342-7.

Arain SR, Ruehlow RM, Uhrich TD, Ebert TJ. The efficacy of dexmedetomidine versus morphine for postoperative analgesia after major inpatient surgery. *Anesth Analg*. 2004;98(1):153-8, table of contents.

Aho MS, Erkola OA, Scheinin H, Lehtinen AM, Korttila KT. Effect of intravenously administered dexmedetomidine on pain after laparoscopic tubal ligation. *Anesth Analg*. 1991;73(2):112-8.

Schnabel A, Meyer-frießem CH, Reichl SU, Zahn PK, Pogatzki-zahn EM. Is intraoperative dexmedetomidine a new option for postoperative pain treatment? A meta-analysis of randomized controlled trials. *Pain*. 2013;154(7):1140-9.

Jeon Y, Jang JS. Dexmedetomidine for the treatment of acute pain from complex regional pain syndrome. *Korean J Anesthesiol*. 2011;61(6):537-8.

Nama S, Meenan DR, Fritz WT. The use of sub-anesthetic intravenous ketamine and adjuvant dexmedetomidine when treating acute pain from CRPS. *Pain Physician*. 2010;13(4):365-8.

Sheehy KA, Finkel JC, Darbari DS, Guerrera MF, Quezado ZM. Dexmedetomidine as an Adjuvant to Analgesic Strategy During Vaso-Occlusive Episodes in Adolescents with Sickle-Cell Disease. *Pain Pract*. 2015;15(8):E90-7.

Phillips WJ, Gadiraju S, Dickey S, Galli R, Lerant AA. Dexmedetomidine relieves pain associated with acute sickle cell crisis. *J Pain Symptom Manage*. 2007;34(4):346-9.

Cimen ZS, Hanci A, Sivrikaya GU, Kilinc LT, Erol MK. Comparison of buccal and nasal dexmedetomidine premedication for pediatric patients. *Paediatr Anaesth*. 2013;23(2):134-8.

Mountain BW, Smithson L, Cramolini M, Wyatt TH, Newman M. Dexmedetomidine as a pediatric anesthetic premedication to reduce anxiety and to deter emergence delirium. *AANA J*. 2011;79(3):219-24.

Sakurai Y, Obata T, Odaka A, Terui K, Tamura M, Miyao H. Buccal administration of dexmedetomidine as a preanesthetic in children. *J Anesth*. 2010;24(1):49-53.

Fallah R, Ferdosian F, Shajari A. Non-Parenteral Medications for Procedural Sedation in Children- A Narrative: Review Article. *Iran J Child Neurol*. 2015;9(3):1-8.

Karaaslan D, Peker TT, Alaca A, et al. Comparison of buccal and intramuscular dexmedetomidine premedication for arthroscopic knee surgery. *J Clin Anesth*. 2006;18(8):589-93.

Mason KP, Lubisch N, Robinson F, Roskos R, Epstein MA. Intramuscular dexmedetomidine: an effective route of sedation preserves background activity for pediatric electroencephalograms. *J Pediatr*. 2012;161(5):927-32.

Ravipati P, Reddy PN, Kumar C, Pradeep P, Pathapati RM, Rajashekar ST. Dexmedetomidine decreases the requirement of ketamine and propofol during burns debridement and dressings. *Indian J Anaesth*. 2014;58(2):138-42.

Sun Y, Liu C, Zhang Y, et al. Low-dose intramuscular dexmedetomidine as premedication: a randomized controlled trial. *Med Sci Monit.* 2014;20:2714-9.

Mason KP, Lubisch NB, Robinson F, Roskos R. Intramuscular dexmedetomidine sedation for pediatric MRI and CT. *AJR Am J Roentgenol.* 2011;197(3):720-5.

Bloor BC, Ward DS, Belleville JP, Maze M. Effects of intravenous dexmedetomidine in humans. II. Hemodynamic changes. *Anesthesiology.* 1992;77(6):1134-42.

McMorrow SP, Abramo TJ. Dexmedetomidine sedation: uses in pediatric procedural sedation outside the operating room. *Pediatr Emerg Care.* 2012;28(3):292-6.

Plambech MZ, Afshari A. Dexmedetomidine in the pediatric population: a review. *Minerva Anesthesiol.* 2015;81(3):320-32.

Dahmani S, Brasher C, Stany I, et al. Premedication with clonidine is superior to benzodiazepines. A meta analysis of published studies. *Acta Anaesthesiol Scand.* 2010;54(4):397-402.

Hayden JC, Breatnach C, Doherty DR, et al. Efficacy of α 2-Agonists for Sedation in Pediatric Critical Care: A Systematic Review. *Pediatr Crit Care Med.* 2016;17(2):e66-75.

Henry K, Harris CR. Deadly ingestions. *Pediatr Clin North Am.* 2006;53(2):293-315.

Seeger DL. Clonidine toxicity revisited. *J Toxicol Clin Toxicol.* 2002;40(2):145-55.

Pertovaara A, Haapalinna A, Sirviö J, Virtanen R. Pharmacological properties, central nervous system effects, and potential therapeutic applications of atipamezole, a selective alpha2-adrenoceptor antagonist. *CNS Drug Rev.* 2005;11(3):273-88.

Sear JW, Higham H. Issues in the perioperative management of the elderly patient with cardiovascular disease. *Drugs Aging.* 2002;19(6):429-51.

Guay DR. Adjunctive agents in the management of chronic pain. *Pharmacotherapy.* 2001;21(9):1070-81.

Bharati S, Pal A, Biswas C, Biswas R. Incidence of cardiac arrest increases with the indiscriminate use of dexmedetomidine: a case series and review of published case reports. *Acta Anaesthesiol Taiwan.* 2011;49(4):165-7.

Horvath JS, Phippard A, Korda A, Henderson-smart DJ, Child A, Tiller DJ. Clonidine hydrochloride--a safe and effective antihypertensive agent in pregnancy. *Obstet Gynecol.* 1985;66(5):634-8.

Rubin PC, Butters L, Clark DM, et al. Placebo-controlled trial of atenolol in treatment of

pregnancy-associated hypertension. *Lancet*. 1983;1(8322):431-4.

Li C, Li Y, Wang K, Kong X. Comparative Evaluation of Remifentanyl and Dexmedetomidine in General Anesthesia for Cesarean Delivery. *Med Sci Monit*. 2015;21:3806-13.

Yousef AA, Salem HA, Moustafa MZ. Effect of mini-dose epidural dexmedetomidine in elective cesarean section using combined spinal-epidural anesthesia: a randomized double-blinded controlled study. *J Anesth*. 2015;29(5):708-14.

Handlogten KS, Sharpe EE, Brost BC, Parney IF, Pasternak JJ. Dexmedetomidine and Mannitol for Awake Craniotomy in a Pregnant Patient. *Anesth Analg*. 2015;120(5):1099-103.

Li Z, Tian M, Zhang CY, et al. A Randomised Controlled Trial to Evaluate the Effectiveness of Intrathecal Bupivacaine Combined with Different Adjuvants (Fentanyl, Clonidine and Dexmedetomidine) in Caesarean Section. *Drug Res (Stuttg)*. 2015;65(11):581-6.

Nie Y, Liu Y, Luo Q, Huang S. Effect of dexmedetomidine combined with sufentanil for post-caesarean section intravenous analgesia: a randomised, placebo-controlled study. *Eur J Anaesthesiol*. 2014;31(4):197-203.

Abu-Halaweh SA, Al Oweidi AK, Abu-Malooh H, et al. Intravenous dexmedetomidine infusion for labour analgesia in patient with preeclampsia. *Eur J Anaesthesiol*. 2009;26(1):86-7.

Sia AT, Kwek K, Yeo GS. The in vitro effects of clonidine and dexmedetomidine on human myometrium. *Int J Obstet Anesth*. 2005;14(2):104-7.

MacMurdo M, Poonyagariyagorn H. 1857: A pregnant pause: the use of dexmedetomidine for agitated delirium in pregnancy. *Crit Care Med*. 2016;44(12 Suppl 1):539.

Pawlik MT, Hansen E, Waldhauser D, Selig C, Kuehnel TS. Clonidine premedication in patients with sleep apnea syndrome: a randomized, double-blind, placebo-controlled study. *Anesth Analg*. 2005;101(5):1374-80.

Sollazzi L, Modesti C, Vitale F, et al. Preinductive use of clonidine and ketamine improves recovery and reduces postoperative pain after bariatric surgery. *Surg Obes Relat Dis*. 2009;5(1):67-71.

Cortínez LI, Anderson BJ, Holford NH, et al. Dexmedetomidine pharmacokinetics in the obese. *Eur J Clin Pharmacol*. 2015;71(12):1501-8.

Ingrande J, Lemmens HJ. Dose adjustment of anaesthetics in the morbidly obese. *Br J Anaesth*. 2010;105 Suppl 1:i16-23.

Calver L, Isbister GK. Dexmedetomidine in the emergency department: assessing safety and

effectiveness in difficult-to-sedate acute behavioural disturbance. *Emerg Med J*. 2012;29(11):915-8.

Gabapentinoids

Field MJ, Cox PJ, Stott E, et al. Identification of the alpha2-delta-1 subunit of voltage-dependent calcium channels as a molecular target for pain mediating the analgesic actions of pregabalin. *Proc Natl Acad Sci USA*. 2006;103(46):17537-42.

Gee NS, Brown JP, Dissanayake VU, Offord J, Thurlow R, Woodruff GN. The novel anticonvulsant drug, gabapentin (Neurontin), binds to the alpha2delta subunit of a calcium channel. *J Biol Chem*. 1996;271(10):5768-76.

Taylor CP. Mechanisms of analgesia by gabapentin and pregabalin--calcium channel alpha2-delta [Cavalpha2-delta] ligands. *Pain*. 2009;142(1-2):13-6.

Kremer M, Salvat E, Muller A, Yalcin I, Barrot M. Antidepressants and gabapentinoids in neuropathic pain: Mechanistic insights. *Neuroscience*. 2016;338:183-206.

Patel R, Dickenson AH. Mechanisms of the gabapentinoids and $\alpha 2 \delta$ -1 calcium channel subunit in neuropathic pain. *Pharmacol Res Perspect*. 2016;4(2):e00205.

Hachad H, Ragueneau-Majlessi I, Levy RH. New antiepileptic drugs: review on drug interactions. *Ther Drug Monit*. 2002;24(1):91-103.

Taylor CP, Angelotti T, Fauman E. Pharmacology and mechanism of action of pregabalin: the calcium channel alpha2-delta (alpha2-delta) subunit as a target for antiepileptic drug discovery. *Epilepsy Res*. 2007;73(2):137-50.

Moore RA, Wiffen PJ, Derry S, Toelle T, Rice AS. Gabapentin for chronic neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev*. 2014;(4):CD007938.

Gabapentin for adults with neuropathic pain: A review of the clinical evidence and guidelines. Canadian Agency for Drugs and Technologies in Health 2014.

Moore RA, Straube S, Wiffen PJ, Derry S, Mcquay HJ. Pregabalin for acute and chronic pain in adults. *Cochrane Database Syst Rev*. 2009;(3):CD007076.

Yan PZ, Butler PM, Kurowski D, Perloff MD. Beyond neuropathic pain: gabapentin use in cancer pain and perioperative pain. *Clin J Pain*. 2014;30(7):613-29.

Fabritius ML, Geisler A, Petersen PL, et al. Gabapentin for post-operative pain management - a systematic review with meta-analyses and trial sequential analyses. *Acta Anaesthesiol Scand*. 2016;60(9):1188-208.

Dahl JB, Nielsen RV, Wetterslev J, et al. Post-operative analgesic effects of paracetamol, NSAIDs, glucocorticoids, gabapentinoids and their combinations: a topical review. *Acta Anaesthesiol Scand*. 2014;58(10):1165-81.

Mishriky BM, Waldron NH, Habib AS. Impact of pregabalin on acute and persistent postoperative pain: a systematic review and meta-analysis. *Br J Anaesth*. 2015;114(1):10-31.

Clarke H, Bonin RP, Orser BA, Englesakis M, Wijeyesundera DN, Katz J. The prevention of chronic postsurgical pain using gabapentin and pregabalin: a combined systematic review and meta-analysis. *Anesth Analg*. 2012;115(2):428-42.

Schmidt PC, Ruchelli G, Mackey SC, Carroll IR. Perioperative gabapentinoids: choice of agent, dose, timing, and effects on chronic postsurgical pain. *Anesthesiology*. 2013;119(5):1215-21.

Butkovic D, Toljan S, Mihovilovic-novak B. Experience with gabapentin for neuropathic pain in adolescents: report of five cases. *Paediatr Anaesth*. 2006;16(3):325-9.

Lauder GR, White MC. Neuropathic pain following multilevel surgery in children with cerebral palsy: a case series and review. *Paediatr Anaesth*. 2005;15(5):412-20.

Friedrichsdorf SJ, Nugent AP. Management of neuropathic pain in children with cancer. *Curr Opin Support Palliat Care*. 2013;7(2):131-8.

Rusy LM, Hainsworth KR, Nelson TJ, et al. Gabapentin use in pediatric spinal fusion patients: a randomized, double-blind, controlled trial. *Anesth Analg*. 2010;110(5):1393-8.

Salman AE, Camkiran A, Oğuz S, Dönmez A. Gabapentin premedication for postoperative analgesia and emergence agitation after sevoflurane anesthesia in pediatric patients. *Agri*. 2013;25(4):163-8.

Mayell A, Srinivasan I, Campbell F, Peliowski A. Analgesic effects of gabapentin after scoliosis surgery in children: a randomized controlled trial. *Paediatr Anaesth*. 2014;24(12):1239-44.

Amani S, Abedinzadeh MR. Effects of Oral Gabapentin, Local Bupivacaine and Intravenous Pethidine on Post Tonsillectomy Pain. *Iran J Otorhinolaryngol*. 2015;27(82):343-8.

Tsai KC, Yang YL, Fan PC. Gabapentin for Postoperative Vomiting in Children Requiring Posterior Fossa Tumor Resection. *Pediatr Neonatol*. 2015;56(5):351-4.

Vondracek P, Oslejskova H, Kepak T, et al. Efficacy of pregabalin in neuropathic pain in paediatric oncological patients. *Eur J Paediatr Neurol*. 2009;13(4):332-6.

Saltık S, Sözen HG, Basgul S, Karatoprak EY, İçağasioğlu A. Pregabalin Treatment of a Patient With Complex Regional Pain Syndrome. *Pediatr Neurol*. 2016;54:88-90.

Tomson T, Landmark CJ, Battino D. Antiepileptic drug treatment in pregnancy: changes in drug disposition and their clinical implications. *Epilepsia*. 2013;54(3):405-14.

Reimers A. New antiepileptic drugs and women. *Seizure*. 2014;23(8):585-91.

Fujii H, Goel A, Bernard N, et al. Pregnancy outcomes following gabapentin use: results of a prospective comparative cohort study. *Neurology*. 2013;80(17):1565-70.

Toth C. Pregabalin: latest safety evidence and clinical implications for the management of neuropathic pain. *Ther Adv Drug Saf*. 2014;5(1):38-56.

Winterfeld U, Merlob P, Baud D, et al. Pregnancy outcome following maternal exposure to pregabalin may call for concern. *Neurology*. 2016;86(24):2251-7.

Lavand'homme PM, Roelants F. Evaluation of pregabalin as an adjuvant to patient-controlled epidural analgesia during late termination of pregnancy. *Anesthesiology*. 2010;113(5):1186-91.

Schifano F. Misuse and abuse of pregabalin and gabapentin: cause for concern?. *CNS Drugs*. 2014;28(6):491-6.

Kapil V, Green JL, Le lait MC, Wood DM, Dargan PI. Misuse of the γ -aminobutyric acid analogues baclofen, gabapentin and pregabalin in the UK. *Br J Clin Pharmacol*. 2014;78(1):190-1.

Stannard C. Misuse of gabapentin and pregabalin: a marker for a more serious malaise?. *Addiction*. 2016;111(10):1699-700.

Chiappini S, Schifano F. A Decade of Gabapentinoid Misuse: An Analysis of the European Medicines Agency's 'Suspected Adverse Drug Reactions' Database. *CNS Drugs*. 2016;30(7):647-54.

Badgaiyan RD. A Novel Perspective on Dopaminergic Processing of Human Addiction. *J Alcohol Drug Depend*. 2013;1(1).

Cai K, Nanga RP, Lamprou L, et al. The impact of gabapentin administration on brain GABA and glutamate concentrations: a $7T$ 1H -MRS study. *Neuropsychopharmacology*. 2012;37(13):2764-71.

Mersfelder TL, Nichols WH. Gabapentin: Abuse, Dependence, and Withdrawal. *Ann Pharmacother*. 2016;50(3):229-33.

Smith RV, Havens JR, Walsh SL. Gabapentin misuse, abuse and diversion: a systematic review. *Addiction*. 2016;111(7):1160-74.

Gahr M, Freudenmann RW, Hiemke C, Kölle MA, Schönfeldt-Iecuona C. Pregabalin abuse and dependence in Germany: results from a database query. *Eur J Clin Pharmacol*. 2013;69(6):1335-42.

Schifano F, D'offizi S, Piccione M, et al. Is there a recreational misuse potential for pregabalin? Analysis of anecdotal online reports in comparison with related gabapentin and clonazepam data. *Psychother Psychosom*. 2011;80(2):118-22.

Schjerning O, Rosenzweig M, Pottegård A, Damkier P, Nielsen J. Abuse Potential of Pregabalin: A Systematic Review. *CNS Drugs*. 2016;30(1):9-25.

Baird CR, Fox P, Colvin LA. Gabapentinoid abuse in order to potentiate the effect of methadone: a survey among substance misusers. *Eur Addict Res*. 2014;20(3):115-8.

Reeves RR, Ladner ME. Potentiation of the effect of buprenorphine/naloxone with gabapentin or quetiapine. *Am J Psychiatry*. 2014;171(6):691.

Sodium Channel Antagonists

Whiteside JB, Wildsmith JA. Developments in local anaesthetic drugs. *Br J Anaesth*. 2001;87(1):27-35.

Nguyen HM, Goldin AL. Sodium channel carboxyl-terminal residue regulates fast inactivation. *J Biol Chem*. 2010;285(12):9077-89.

Catterall WA, Swanson TM. Structural Basis for Pharmacology of Voltage-Gated Sodium and Calcium Channels. *Mol Pharmacol*. 2015;88(1):141-50.

Lipkind GM, Fozzard HA. Molecular modeling of local anesthetic drug binding by voltage-gated sodium channels. *Mol Pharmacol*. 2005;68(6):1611-22.

Hille B. Local anesthetics: hydrophilic and hydrophobic pathways for the drug-receptor reaction. *J Gen Physiol*. 1977;69(4):497-515.

Eipe N, Gupta S, Penning J. Intravenous lidocaine for acute pain: An evidence-based clinical update. *BJA Education* 2016:1-7.

Vanstone RJ, Rockett M. Use of atypical analgesics by intravenous infusion (IV) for acute pain: evidence base for lidocaine, ketamine and magnesium. *Anesthesia & Intensive Care Medicine* 2016; 17(9): 460-463.

Lauretti GR. Mechanisms of analgesia of intravenous lidocaine. *Rev Bras Anesthesiol*. 2008;58(3):280-6.

Benowitz NL, Meister W. Clinical pharmacokinetics of lignocaine. *Clin Pharmacokinet*. 1978;3(3):177-201.

Link MS, Berkow LC, Kudenchuk PJ, et al. Part 7: Adult Advanced Cardiovascular Life Support: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2015;132(18 Suppl 2):S444-64.

McLure HA, Rubin AP. Review of local anaesthetic agents. *Minerva Anesthesiol*. 2005;71(3):59-74.

Welling LR, Watson WA. The emergency department treatment of dyspepsia with antacids and oral lidocaine. *Ann Emerg Med*. 1990;19(7):785-8.

Vilke GM, Jin A, Davis DP, Chan TC. Prospective randomized study of viscous lidocaine versus benzocaine in a GI cocktail for dyspepsia. *J Emerg Med*. 2004;27(1):7-9.

Hopper SM, McCarthy M, Tancharoen C, Lee KJ, Davidson A, Babl FE. Topical lidocaine to improve oral intake in children with painful infectious mouth ulcers: a blinded, randomized, placebo-controlled trial. *Ann Emerg Med*. 2014;63(3):292-9.

Ali S, McGrath T, Drendel AL. An Evidence-Based Approach to Minimizing Acute Procedural Pain in the Emergency Department and Beyond. *Pediatr Emerg Care*. 2016;32(1):36-42.

Williams H, Carlton E. Towards evidence based emergency medicine: best BETs from the Manchester Royal Infirmary. BET 1: Topical lignocaine patches in traumatic rib fractures. *Emerg Med J*. 2015;32(4):333-4.

Prabhakar H, Rath S, Kalaivani M, Bhandari N. Adrenaline with lidocaine for digital nerve blocks. *Cochrane Database Syst Rev*. 2015;(3):CD010645.

Mohr B. Safety and effectiveness of intravenous regional anesthesia (Bier block) for outpatient management of forearm trauma. *CJEM*. 2006;8(4):247-50.

Grabinsky A, Sharar SR. Regional anesthesia for acute traumatic injuries in the emergency room. *Expert Rev Neurother*. 2009;9(11):1677-90.

Blasier RD, White R. Intravenous regional anesthesia for management of children's extremity fractures in the emergency department. *Pediatr Emerg Care*. 1996;12(6):404-6.

Tirado A, Wu T, Noble VE, et al. Ultrasound-guided procedures in the emergency department-diagnostic and therapeutic asset. *Emerg Med Clin North Am*. 2013;31(1):117-49.

Liebmann O, Price D, Mills C, et al. Feasibility of forearm ultrasonography-guided nerve blocks of the radial, ulnar, and median nerves for hand procedures in the emergency department. *Ann Emerg Med*. 2006;48(5):558-62.

Blaivas M, Adhikari S, Lander L. A prospective comparison of procedural sedation and ultrasound-guided interscalene nerve block for shoulder reduction in the emergency department. *Acad Emerg Med*. 2011;18(9):922-7.

Beaudoin FL, Nagdev A, Merchant RC, Becker BM. Ultrasound-guided femoral nerve blocks

in elderly patients with hip fractures. *Am J Emerg Med.* 2010;28(1):76-81.

Reid N, Stella J, Ryan M, Ragg M. Use of ultrasound to facilitate accurate femoral nerve block in the emergency department. *Emerg Med Australas.* 2009;21(2):124-30.

Stone MB, Wang R, Price DD. Ultrasound-guided supraclavicular brachial plexus nerve block vs procedural sedation for the treatment of upper extremity emergencies. *Am J Emerg Med.* 2008;26(6):706-10.

Herring AA, Stone MB, Frenkel O, Chipman A, Nagdev AD. The ultrasound-guided superficial cervical plexus block for anesthesia and analgesia in emergency care settings. *Am J Emerg Med.* 2012;30(7):1263-7.

Beaudoin FL, Haran JP, Liebmann O. A comparison of ultrasound-guided three-in-one femoral nerve block versus parenteral opioids alone for analgesia in emergency department patients with hip fractures: a randomized controlled trial. *Acad Emerg Med.* 2013;20(6):584-91.

Haines L, Dickman E, Ayvazyan S, et al. Ultrasound-guided fascia iliaca compartment block for hip fractures in the emergency department. *J Emerg Med.* 2012;43(4):692-7.

Guay J, Suresh S, Kopp S. The use of ultrasound guidance for perioperative neuraxial and peripheral nerve blocks in children. *Cochrane Database Syst Rev.* 2016;2:CD011436.

Fitch RW, Kuhn JE. Intraarticular lidocaine versus intravenous procedural sedation with narcotics and benzodiazepines for reduction of the dislocated shoulder: a systematic review. *Acad Emerg Med.* 2008;15(8):703-8.

Wakai A, O'Sullivan R, McCabe A. Intra-articular lignocaine versus intravenous analgesia with or without sedation for manual reduction of acute anterior shoulder dislocation in adults. *Cochrane Database Syst Rev.* 2011;(4):CD004919.

Jiang N, Hu YJ, Zhang KR, Zhang S, Bin Y. Intra-articular lidocaine versus intravenous analgesia and sedation for manual closed reduction of acute anterior shoulder dislocation: an updated meta-analysis. *J Clin Anesth.* 2014;26(5):350-9.

White BJ, Walsh M, Egol KA, Tejwani NC. Intra-articular block compared with conscious sedation for closed reduction of ankle fracture-dislocations. A prospective randomized trial. *J Bone Joint Surg Am.* 2008;90(4):731-4.

Vahidi E, Shakoor D, Aghaie meybodi M, Saeedi M. Comparison of intravenous lidocaine versus morphine in alleviating pain in patients with critical limb ischaemia. *Emerg Med J.* 2015;32(7):516-9.

Tanen DA, Shimada M, Danish DC, Dos santos F, Makela M, Riffenburgh RH. Intravenous

lidocaine for the emergency department treatment of acute radicular low back pain, a randomized controlled trial. *J Emerg Med.* 2014;47(1):119-24.

Reutens DC, Fatovich DM, Stewart-wynne EG, Prentice DA. Is intravenous lidocaine clinically effective in acute migraine?. *Cephalalgia.* 1991;11(6):245-7.

Bell R, Montoya D, Shuaib A, Lee MA. A comparative trial of three agents in the treatment of acute migraine headache. *Ann Emerg Med.* 1990;19(10):1079-82.

Soleimanpour H, Hassanzadeh K, Mohammadi DA, Vaezi H, Esfanjani RM. Parenteral lidocaine for treatment of intractable renal colic: a case series. *J Med Case Rep.* 2011;5:256.

Soleimanpour H, Hassanzadeh K, Vaezi H, Golzari SE, Esfanjani RM, Soleimanpour M. Effectiveness of intravenous lidocaine versus intravenous morphine for patients with renal colic in the emergency department. *BMC Urol.* 2012;12:13.

Sin B, Effendi M, Bjork C, Punnapuza S. The Use of Intravenous Lidocaine for Renal Colic in the Emergency Department. *Ann Pharmacother.* 2016;50(3):242.

Firouzian A, Alipour A, Rashidian Dezfouli H, et al. Does lidocaine as an adjuvant to morphine improve pain relief in patients presenting to the ED with acute renal colic? A double-blind, randomized controlled trial. *Am J Emerg Med.* 2016;34(3):443-8.

Baranowski AP, De Courcey J, Bonello E. A trial of intravenous lidocaine on the pain and allodynia of postherpetic neuralgia. *J Pain Symptom Manage.* 1999;17(6):429-33.

Neal JM, Bernardis CM, Hadzic A, et al. ASRA Practice Advisory on Neurologic Complications in Regional Anesthesia and Pain Medicine. *Reg Anesth Pain Med.* 2008;33(5):404-15.

Welch MB, Brummett CM, Welch TD, et al. Perioperative peripheral nerve injuries: a retrospective study of 380,680 cases during a 10-year period at a single institution. *Anesthesiology.* 2009;111(3):490-7.

Faccenda KA, Finucane BT. Complications of regional anaesthesia Incidence and prevention. *Drug Saf.* 2001;24(6):413-42.

Freedman JM, Li DK, Drasner K, Jaskela MC, Larsen B, Wi S. Transient neurologic symptoms after spinal anesthesia: an epidemiologic study of 1,863 patients. *Anesthesiology.* 1998;89(3):633-41.

Pollock JE. Transient neurologic symptoms: etiology, risk factors, and management. *Reg Anesth Pain Med.* 2002;27(6):581-6.

Zink W, Graf BM. Local anesthetic myotoxicity. *Reg Anesth Pain Med.* 2004;29(4):333-40.

Tanawuttiwat T, Thisayakorn P, Viles-Gonzalez JF. LAST (local anesthetic systemic toxicity) but not least: systemic lidocaine toxicity during cardiac intervention. *J Invasive Cardiol.* 2014;26(1):E13-5.

Dorf E, Kuntz AF, Kelsey J, Holstege CP. Lidocaine-induced altered mental status and seizure after hematoma block. *J Emerg Med.* 2006;31(3):251-3.

Drasner K. Local anesthetic neurotoxicity: clinical injury and strategies that may minimize risk. *Reg Anesth Pain Med.* 2002;27(6):576-80.

Covino BG. Pharmacology of local anesthetic agents. *Ration Drug Ther.* 1987;21(8):1-9.

Alfano SN, Leicht MJ, Skiendzielewski JJ. Lidocaine toxicity following subcutaneous administration. *Ann Emerg Med.* 1984;13(6):465-7.

Gosselin S, Hoegberg LC, Hoffman RS, et al. Evidence-based recommendations on the use of intravenous lipid emulsion therapy in poisoning *Clin Toxicol (Phila).* 2016;1-25. [Epub ahead of print]

Neal JM, Bernardis CM, Butterworth JF, et al. ASRA practice advisory on local anesthetic systemic toxicity. *Reg Anesth Pain Med.* 2010;35(2):152-61.

Guay J. Methemoglobinemia related to local anesthetics: a summary of 242 episodes. *Anesth Analg.* 2009;108(3):837-45.

Barash M, Reich KA, Rademaker D. Lidocaine-induced methemoglobinemia: a clinical reminder. *J Am Osteopath Assoc.* 2015;115(2):94-8.

Kennedy RM, Luhmann JD. Pharmacological management of pain and anxiety during emergency procedures in children. *Paediatr Drugs.* 2001;3(5):337-54.

Blasier RD, White R. Intravenous regional anesthesia for management of children's extremity fractures in the emergency department. *Pediatr Emerg Care.* 1996;12(6):404-6.

Barnett P. Alternatives to sedation for painful procedures. *Pediatr Emerg Care.* 2009;25(6):415-9.

Breslin K, Boniface K, Cohen J. Ultrasound-guided intra-articular lidocaine block for reduction of anterior shoulder dislocation in the pediatric emergency department. *Pediatr Emerg Care.* 2014;30(3):217-20.

Marin JR, Abo AM, Arroyo AC, et al. Pediatric emergency medicine point-of-care ultrasound: summary of the evidence. *Crit Ultrasound J.* 2016;8(1):16. [Epub ahead of print]

Constantine E, Tsze DS, Machan JT, Ebersson CP, Linakis JG, Steele DW. Evaluating the

hematoma block as an adjunct to procedural sedation for closed reduction of distal forearm fractures. *Pediatr Emerg Care*. 2014;30(7):474-8.

Constantine E, Steele DW, Ebersson C, Boutis K, Amanullah S, Linakis JG. The use of local anesthetic techniques for closed forearm fracture reduction in children: a survey of academic pediatric emergency departments. *Pediatr Emerg Care*. 2007;23(4):209-11.

Baker MD, Gullett JP. Ultrasound-Guided Femoral Nerve Blocks. *Pediatr Emerg Care*. 2015;31(12):864-8.

Neubrand TL, Roswell K, Deakyne S, Kocher K, Wathen J. Fascia iliaca compartment nerve block versus systemic pain control for acute femur fractures in the pediatric emergency department. *Pediatr Emerg Care*. 2014;30(7):469-73.

Turner AL, Stevenson MD, Cross KP. Impact of ultrasound-guided femoral nerve blocks in the pediatric emergency department. *Pediatr Emerg Care*. 2014;30(4):227-9.

Black KJ, Bevan CA, Murphy NG, Howard JJ. Nerve blocks for initial pain management of femoral fractures in children. *Cochrane Database Syst Rev*. 2013;(12):CD009587.

Bretholz A, Doan Q, Cheng A, Lauder G. A presurvey and postsurvey of a web- and simulation-based course of ultrasound-guided nerve blocks for pediatric emergency medicine. *Pediatr Emerg Care*. 2012;28(6):506-9.

Bolte RG, Stevens PM, Scott SM, Schunk JE. Mini-dose Bier block intravenous regional anesthesia in the emergency department treatment of pediatric upper-extremity injuries. *J Pediatr Orthop*. 1994;14(4):534-7.

Rosenberg PH, Veering BT, Urmey WF. Maximum recommended doses of local anesthetics: a multifactorial concept. *Reg Anesth Pain Med*. 2004;29(6):564-75.

American Geriatrics Society Panel on Pharmacological Management of Persistent Pain in Older Persons. Pharmacological management of persistent pain in older persons. *J Am Geriatr Soc*. 2009;57(8):1331-46.

Halaszynski TM. Pain management in the elderly and cognitively impaired patient: the role of regional anesthesia and analgesia. *Curr Opin Anaesthesiol*. 2009;22(5):594-9.

Beaudoin FL, Nagdev A, Merchant RC, Becker BM. Ultrasound-guided femoral nerve blocks in elderly patients with hip fractures. *Am J Emerg Med*. 2010;28(1):76-81.

Beaudoin FL, Haran JP, Liebmann O. A comparison of ultrasound-guided three-in-one femoral nerve block versus parenteral opioids alone for analgesia in emergency department patients with hip fractures: a randomized controlled trial. *Acad Emerg Med*. 2013;20(6):584-91.

Ritcey B, Pageau P, Woo MY, Perry JJ. Regional Nerve Blocks For Hip and Femoral Neck Fractures in the Emergency Department: A Systematic Review. *CJEM*. 2016;18(1):37-47.

Santos AC, Pedersen H, Harmon TW, et al. Does pregnancy alter the systemic toxicity of local anesthetics?. *Anesthesiology*. 1989;70(6):991-5.

Moller RA, Datta S, Strichartz GR. Beta-estradiol acutely potentiates the depression of cardiac excitability by lidocaine and bupivacaine. *J Cardiovasc Pharmacol*. 1999;34(5):718-27.

Moller RA, Datta S, Fox J, Johnson M, Covino BG. Effects of progesterone on the cardiac electrophysiologic action of bupivacaine and lidocaine. *Anesthesiology*. 1992;76(4):604-8.

Finster M. Toxicity of local anesthetics in the fetus and the newborn. *Bull N Y Acad Med*. 1976;52(2):222-5.

Govindappagari S, Grossman TB, Dayal AK, Grosberg BM, Vollbracht S, Robbins MS. Peripheral nerve blocks in the treatment of migraine in pregnancy. *Obstet Gynecol*. 2014;124(6):1169-74.

Peleg R, Gohar J, Koretz M, Peleg A. Abdominal wall pain in pregnant women caused by thoracic lateral cutaneous nerve entrapment. *Eur J Obstet Gynecol Reprod Biol*. 1997;74(2):169-71.

Shenkman Z, Shir Y, Brodsky JB. Perioperative management of the obese patient. *Br J Anaesth*. 1993;70(3):349-59.

Lottia S. Anaesthesia and Morbid Obesity. *Contin Educ Anaesth Crit Care Pain* 2008; 8(5):151-156.

Ingrande J, Lemmens HJ. Dose adjustment of anaesthetics in the morbidly obese. *Br J Anaesth*. 2010;105 Suppl 1:i16-23.

Roberson DP, Binshtok AM, Blasl F, Bean BP, Woolf CJ. Targeting of sodium channel blockers into nociceptors to produce long-duration analgesia: a systematic study and review. *Br J Pharmacol*. 2011;164(1):48-58.

Bagal SK, Chapman ML, Marron BE, Prime R, Storer RI, Swain NA. Recent progress in sodium channel modulators for pain. *Bioorg Med Chem Lett*. 2014;24(16):3690-9.

Bagal SK, Marron BE, Owen RM, Storer RI, Swain NA. Voltage gated sodium channels as drug discovery targets. *Channels (Austin)*. 2015;9(6):360-6.

Verdu B, Decosterd I, Buclin T, Stiefel F, Berney A. Antidepressants for the treatment of

chronic pain. *Drugs*. 2008;68(18):2611-32.

Jackson JL, Shimeall W, Sessums L, et al. Tricyclic antidepressants and headaches: systematic review and meta-analysis. *BMJ*. 2010;341:c5222.

Wong K, Phelan R, Kalso E, et al. Antidepressant drugs for prevention of acute and chronic postsurgical pain: early evidence and recommended future directions. *Anesthesiology*. 2014;121(3):591-608.

Joslin J, Worthing R, Ladbroke M, Mularella J. Amitriptyline use for acute pain in remote environments. *Wilderness Environ Med*. 2014;25(4):488-9.

Sawynok J, Esser MJ, Reid AR. Antidepressants as analgesics: an overview of central and peripheral mechanisms of action. *J Psychiatry Neurosci*. 2001;26(1):21-9.

Kremer M, Salvat E, Muller A, Yalcin I, Barrot M. Antidepressants and gabapentinoids in neuropathic pain: Mechanistic insights. *Neuroscience*. 2016;338:183-206.

Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain - United States, 2016. *MMWR Recomm Rep*. 2016;65(1):1-49.